

# POST-OPERATIVE INSTRUCTIONS TO CARE FOR YOUR CHILD

Your child has received one or more of the following circled items. Please follow the instructions below for his/her proper care.

Feel free to call: **Dr. Joni Price** (256-329-8401)

<p><b>Sedative</b></p>	<ul style="list-style-type: none"> <li>◆Your child will probably be sleepy or acting groggy today due to receiving a sedative. Don't be surprised if your child wants to take a nap for several hours, just make sure your child rests on his/her side or back and is closely monitored by an adult for the rest of the day. Activities should be limited to watching TV, playing quietly with toys inside and resting.</li> <li>◆Your child should not return to school or daycare today, he/she needs to be under close parental supervision. If work related obligations necessitate that your child return to school/daycare, please inform the teacher/caregiver that your child is to be monitored closely.</li> <li>◆<b>Do not plan or permit activities for your child after treatment.</b> Your child's activities should be <b>limited &amp; supervised</b> today. This means no running, bicycle riding, roller-skating, swinging on a swing, swimming at the pool or playing at the playground.</li> <li>◆Your child may experience some nausea. If so, please allow only sips of clear liquids (Gatorade &amp; Pedialite) followed by softer foods. Please do not give heavy foods (like hamburgers or pizza) to start with.</li> </ul>
<p><b>Local Anesthetic</b></p>	<ul style="list-style-type: none"> <li>◆Your child has been given an injection of local anesthetic to numb his/her teeth in order for us to work on them. Your child's mouth (tooth, lip &amp; tongue) will be sleeping for <b>approximately 2 to 3 hours</b>. It is imperative that you watch your child and have him/her bite on the cotton roll provided to prevent damage to his/her oral cavity. If your child is <b>not biting</b> on the cotton roll he/she may bite the tongue, cheek or lip causing <b>serious damage</b> or profound <b>pain</b> later on.</li> <li>◆If your child falls asleep you may remove the cotton roll to prevent him/her from swallowing or choking on it.</li> <li>◆Some children become <b>very upset</b> (even crying hysterically) and complain of excruciating pain when they realize their mouth feels "different." Please <b>do not be alarmed!</b> Many children are unfamiliar with this numb sensation and associate this with pain. Reassure your child that the "funny feeling" will go away in approximately 2 hours.</li> </ul>
<p><b>Cleaning</b></p>	<ul style="list-style-type: none"> <li>◆Your child has received a dental a cleaning which included the application of fluoride. The fluoride is applied to the teeth with a toothbrush after the cleaning in an effort to prevent cavities from developing. Because of this it is important that you not allow your child to eat or drink anything for 20 minutes to enable the fluoride to "soak in." The application of fluoride is recommended every 6 months.</li> </ul>
<p><b>Sealant</b></p>	<ul style="list-style-type: none"> <li>◆A sealant has been placed on one or more of your child's teeth in an effort to prevent him/her from developing a cavity on the top surface of the tooth. Your child must avoid eating ice or chewing on hard sticky candies, which can dislodge or chip the sealant, rendering it ineffective. Please also be advised that sealants do not protect the surfaces in-between teeth so your child must still floss to prevent this type of cavity from developing between the teeth.</li> </ul>
<p><b>Filling</b></p>	<ul style="list-style-type: none"> <li>◆Most children usually adapt and adjust very well to new dental work. However, if some fillings are a bit deep due to the extent of the decay, they may complain of some discomfort. Do not be alarmed! Sometimes it takes a few weeks for the tooth to settle down and feel comfortable. Your child may have received a silver or white (composite) colored filling.</li> </ul>
<p><b>Stainless Steel Crown (Cap)</b></p>	<ul style="list-style-type: none"> <li>◆If your child has received a crown/cap the gum tissue surrounding the tooth may appear to be bleeding. This is normal! When preparing your child's tooth for a crown, it is necessary to fit the crown closely to the gum tissue, resulting in some hemorrhaging and bruising of the tissue. Following placement of the crowns some parents will notice a purple or gray color around the gums; this is also normal and will subside.</li> <li>◆Please do not allow your child to eat sticky or caramel type candy as this may dislodge your child's crown. This applies for as long as the crown is in your child's mouth.</li> <li>◆If your child's crown does come off please save the crown and call the office so it may be recemented.</li> </ul>
<p><b>Pulpotomy (Nerve Treatment)</b></p>	<ul style="list-style-type: none"> <li>◆Please note that children <b>seldom</b> complain of pain after this procedure is performed. When your child receives a nerve treatment it is usually <b>not</b> necessary to prescribe any pain medication. Children that do experience discomfort usually do fine with Tylenol or Ibuprofen (follow directions on the bottle).</li> </ul>
<p><b>Extraction</b></p>	<ul style="list-style-type: none"> <li>◆ You may resume brushing tonight, brushing gently in the site of the extraction.</li> <li>◆<b>Starting tomorrow</b> you may use a warm salt water rinses to keep the extraction site clean. Simply place 1 to 2 teaspoons of salt in a warm cup of water - stir and then have your child swish by mouth 3 to 6 times daily, or at least after each meal or snack.</li> <li>◆ If your child is too young to swish, simply dip a wash cloth in a the mixture and swab the extraction site well after each feeding. This area should heal up in approximately 2 to 3 weeks.</li> <li>◆Soft foods are usually tolerated the best for the first few days following removal of the tooth/teeth.</li> <li>◆Do not allow your child to suck from a straw or drink carbonated beverages for 48 hours.</li> </ul>
<p><b>Pain</b></p>	<ul style="list-style-type: none"> <li>◆If your child appears to be in any pain or discomfort give Tylenol, following <b>the instructions on the back of the bottle</b>. If the doctor feels that your child will need a medication stronger, then one will be prescribed. Please note that most children, even with extensive dental work, do very well post-operatively with regular Tylenol Elixer or Ibuprofen. Do not hesitate to call the doctor if you have any questions regarding medications.</li> </ul>
<p><b>What to Eat</b></p>	<ul style="list-style-type: none"> <li>◆Your child should have his/her diet limited to soft foods for 2 hours following the dental procedure to prevent biting the lip or tongue while numb. The following foods are acceptable: milk shake, ice cream, pudding, jello, soup, mashed potatoes, noodles, yogurt or any other foods of this consistency.</li> </ul>